Bound state eigenfunctions need to vanish faster than $|x|^{-3/2}$

Zafar Ahmed

*Nuclear Physics Division, Bhabha Atomic Research Centre, Mumbai 400 085, India**

(Dated: June 6, 2016)

Abstract

In quantum mechanics students are taught to practice that eigenfunction of a physical bound state must be continuous and vanishing asymptotically so that it is normalizable in $x \in (-\infty, \infty)$. Here we caution that such states may also give rise to infinite uncertainty in position ($\Delta x = \infty$), whereas Δp remains finite. Such states may be called loosely bound and spatially extended states that may be avoided by an additional condition that the eigenfunction vanishes asymptotically faster than $|x|^{-3/2}$.

arXiv:1605.05902v2 [quant-ph] 3 Jun 2016

*Electronic address: 1:zahmed@barc.gov.in

The bound state is an essential topic in the curriculum of introductory quantum mechanics [1,2]. Bound state eigenfunctions $\psi_n(x)$ are solutions of the Schrödinger (1926) equation [1,2]

$$\frac{d^2\psi}{dx^2} + \frac{2m}{\hbar^2}[E - V(x)]\psi(x) = 0, \tag{1}$$

for some potential $V(x)$ for specific allowed values of energies $E = E_n$ that obey Dirichlet boundary condition

$$\psi_n(\pm\infty) = 0. \tag{2}$$

Energies E_n are called eigenvalues leading to quantization of energy. This quantization was first hypothesized by Planck(1900) for the explanation of the black body radiation and justified much later by Bohr and Sommerfeld (1919) [1,2] semi-classically for $V(x) = \frac{1}{2}m\omega^2x^2$ and other potential profiles with a single minimum called potential wells.

As $|\psi(x)|^2$ plays the role of quantal probability, $\psi(x)$ is taught to be continuous function on $x \in (-\infty, \infty)$ so that $\psi(x)$ is square integrable (normalizable)

$$\int_{-\infty}^{\infty} |\psi(x)|^2 dx = 1. \tag{3}$$

The additional condition of differentiability of $\psi(x)$ at each and every point of the domain ensures the continuity of the momentum: $-i\hbar\frac{d\psi(x)}{dx}$ unless there is a Dirac delta function in the potential [2]. When the potential contains a Dirac delta function e.g, $g\delta(x-a)$, the eigenfunction is allowed to become non-differentiable at $x = a$ wherein the left and right derivatives are finite but unequal at $x = a$ [3].

Further the uncertainty relation of Heisenberg (1927) [1,2]

$$U = \Delta x \Delta p \geq \frac{\hbar}{2}, \quad \Delta A = \sqrt{< A^2 > - < A >^2}, \quad < A >= \frac{\int_{-\infty}^{\infty} \psi^*(x)\ A\ \psi(x)dx}{\int_{-\infty}^{\infty} \psi^*(x)\psi(x)dx}, \tag{4}$$

requires the uncertainties Δx and Δp to be finite for physically relevant cases. An interesting collection of uncertainty products for ground states of several solvable models may be seen in Ref.[4]. It may be mentioned that nowadays, the terms uncertainty relation and uncertainty principle are classified clearly. The term uncertainty principle is used when an inequality expresses the relationship between measurement error and disturbance. For a recent and a very interesting discussion on this, one may see Ref. [5].

A definite integral $\int_a^b f(x)dx$ is called proper if it represents area under the curve $y = f(x)$ from $x = a$ to $x = b$. This in turn requires $f(x)$ to be continuous or piece-wise continuous in

the closed interval $[a, b]$. On the other hand, an improper [5] integral may not connect well to the area under the curve e.g., $\int_0^1 x^{-1/2}dx = 2$. One way an integral becomes improper is when one or both of its limits are infinite. Improper integrals may or may not be convergent (finite). It is trivial yet important to see that the integral $\int_1^\infty x^{-\beta}dx$ is convergent only if $\beta > 1$. This last point is crucial in determining the asymptotic behavior of $\psi(x)$ such that $< x^2 >$ is finite, see below.

The finiteness of Δx requires $< x^2 >$ to be finite, thus we demand

$$\int_{-\infty}^{\infty} x^2|\psi(x)|^2dx < \infty \Rightarrow |\psi(x)| < |x|^{-3/2}, \quad |x| \sim \infty. \tag{5}$$

On the other hand for $< p >$ and $< p^2 >$ to be finite, $|\psi(x)| \sim 1/x^\epsilon, \epsilon > 0$ would suffice well. Thus, the condition (5) would ensure finiteness of the uncertainty product U [5]. In this paper, we point out this additional condition (5) on the eigenfunction for a bound state.

One can write (1) in an interesting form as

$$V(x) - E_0 = \frac{\hbar^2}{2m}\frac{1}{\psi_0(x)}\frac{d^2\psi_0(x)}{dx^2}. \tag{6}$$

Here we take $2m = 1 = \hbar$ which gives $\frac{2m}{\hbar^2} = 1$. It may be noted that for particle of mass 4 times that of an electron, the approximate value of $\frac{2m}{\hbar^2}$ is $1(eVA^{0\ 2})^{-1}$ when the mass and energy are in eV and length is in A^0. If we put the well known ground state of harmonic oscillator $\psi_0(x) = e^{-x^2/2}$ in (6), we get $V(x) - E_0 = -1 + x^2$. One may interpret that $V(x) = x^2$ and $E_0 = 1$ (or $V(x) = x^2 - 1$ and $E_0 = 0$). This $V(x)$ is the harmonic oscillator potential (See Fig. 1(a)) $V(x) = \frac{1}{2}m\omega^2x^2$, where $\hbar\omega = 2eV$ and $E_0 = 1eV$. The ground state $\psi_0(x) = e^{-x^2/2}$ vanishes faster than $|x|^{-3/2}$ because as $x \to \infty$, $|x|^{3/2}e^{-x^2/2} \to 0 < 1$. So the uncertainty product U_0 for this ground state turns out to be finite as $U_0 = \hbar/2$ [1,2], which equals the least bound on U (4) for any physical eigenstate.

Particularly, in this special case of Harmonic oscillator constructing the raising and lowering operators [1,2] one can find all eigenvalues and corresponding eigenfunctions of $V(x)$. This is a rare feature of a potential.

But if we put $\psi_0(x) = e^{-x^4}$ in (6), we get $V(x) = 16x^6 - 12x^2$ and $E_0 = 0$. See fig. 1(b) this $V(x)$ is a double well and $E_0 = 0$ is the ground state of this potential that possesses infinitely many bound states. One can not construct the raising and lowering operators to obtain the rest of the spectrum of this double well potential. If required the other bound states are obtained by numerically [7] solving the Schrödinger equation for this potential.



FIG. 1: Depiction of the normalized ground states $\psi_0(x)$ (dashed) and the corresponding potentials $V(x)$ (solid) : (a): $\psi_0(x) = (\pi)^{-1/4}\ e^{-x^2/2}$, (b): $\psi_0(x) = [2^{3/4}\Gamma(5/4)]^{-1/2}\ e^{-x^4}$, (c): $\psi_0(x) = 1/\sqrt{\pi(1+x^2)}$. Notice that in the last case: part(c), $V(x)$ (16) vanishes as $\sim 1/x^2$ and its ground state $\psi_0(x)$ (10) vanishes as $\sim 1/x$ [much less rapidly as compared to the previous two ground states (a,b)], hence it represents an extended state.

Once again $\psi_0(x) = e^{-x^4}$ vanishes asymptotically much rapidly as $x \to \infty$, $|x|^{3/2}e^{-x^4} \to 0 < 1$ rendering the integrals for $< x >$ and $< x^2 >$ as finite. Next, $< x >$ vanishes as there is an odd integrand within the symmetric limit of integration. Similarly, $< p >= 0$ in general for bound states as they do not carry any momentum. We can find $< x^2 >$

$$< \psi_0(x)|x^2|\psi_0(x) >= \frac{\int_{-\infty}^{\infty} x^2 e^{-2x^4} dx}{\int_{-\infty}^{\infty} e^{-2x^4} dx} = \frac{\Gamma(3/4)}{4\sqrt{2}\Gamma(5/4)} \tag{7}$$

and

$$< \psi_0(x)|p^2|\psi_0(x) >=< p\psi_0(x)|p\psi_0(x) >= 16\hbar^2 \frac{\int_{-\infty}^{\infty} x^6 e^{-2x^4} dx}{\int_{-\infty}^{\infty} e^{-2x^4} dx} = \frac{\sqrt{2}\Gamma(7/4)}{\Gamma(5/4)}. \tag{8}$$

In Eqs. (7,8), we have converted the integrals to the standard Gamma functions as $\Gamma(z) = \int_0^{\infty} t^{z-1}e^{-t}dt$ [8]. Using (7,8) in (4), we get the uncertainty product for ground state as

$$U_0 = \hbar \frac{\sqrt{\Gamma(3/4)\Gamma(7/4)}}{2\Gamma(5/4)} = 0.5854\ \hbar, \tag{9}$$

which is more than $\hbar/2$ as per the uncertainty principle.

Similarly, the ground state $\psi_0(x) = \sqrt{\frac{2}{\pi}}\frac{1}{1+x^2}$ vanishes a little faster than $|x|^{-3/2}$ to have a well known value for $U_0 = \hbar/\sqrt{2}$ [3].

Now let us take a continuous, differentiable and normalized state as

$$\psi_0(x) = \frac{1}{\sqrt{\pi(1+x^2)}}, \tag{10}$$

which behaves as $1/x$ as $x \sim \infty$. For this state

$$< x >= \frac{1}{\pi}\int_{-\infty}^{\infty}\frac{x}{1+x^2}dx, \quad < x^2 >= \frac{1}{\pi}\int_{-\infty}^{\infty}\frac{x^2}{1+x^2}dx, \tag{11}$$

so both of these integrals (11) are improper. Under the condition that the integration range becomes infinite while maintaining the left-right symmetry of the integration range, we get

$$\Delta x = \infty. \tag{12}$$

On the other hand the integral in

$$< p >= \frac{i\hbar}{\pi}\int_{-\infty}^{\infty}\frac{x}{(1+x^2)^2}dx, \tag{13}$$

is improper but convergent because its integrand vanishes as $|x|^{-3}(\beta > 1)$ asymptotically and then it vanishes as its integrand is an odd function. Similarly $< p^2 >$ is improper but convergent because its integrand vanishes as $|x|^{-4}(\beta > 1)$ asymptotically and it can be evaluated as shown below

$$< p^2 >=< p\psi_0|p\psi_0 >= \frac{\hbar^2}{\pi}\int_{-\infty}^{\infty}\frac{x^2}{(1+x^2)^3}dx = \frac{\hbar^2}{4\pi}\int_0^{\pi/2}[1-\cos 4\theta]\, d\theta = \frac{\hbar^2}{8}. \tag{14}$$

In the integral above, we have used the substitution $x = \tan\theta$, finally we get a finite value of Δp as

$$\Delta p = \frac{\hbar}{2\sqrt{2}}. \tag{15}$$

Eventually, Eqs. (12,15) render the uncertainty product as infinite.

Now let us find out from Eq. (6) the potential giving rise to such a ground state (10), we find

$$V(x) = \frac{2x^2-1}{(1+x^2)^2}. \tag{16}$$

This potential is depicted in Fig. 1(c) and the state (10) is its ground state with eigenvalue $E_0 = 0$. No state can exist below $E = 0$ as it is a node less (ground state). For energies $E > 0$ there are scattering states, there may also exist a metastable (quasi bound, resonant state [9]). Hence the state (10) is the only bound state of this potential (16).

The examples of the ground states $\psi_0(x)$ rendering a finite uncertainty products presented here are instructive which bring out the crucial role of faster asymptotic convergence of $\psi_0(x)$ than $|x|^{-3/2}$. On the other hand the example (10), demonstrates our point that if an

eigenstate of a bound state does not vanish faster than $|x|^{-3/2}$, despite being continuous, differentiable and normalizable it would yield the uncertainty product U as infinite.

Similarly, for 3 dimensional central potentials [1,2] in the radial Schrödinger equation, where $r \in (0, \infty)$ and $\psi(r) = \frac{u(r)}{r}$, $u(r)$ needs to vanish faster than $r^{-3/2}$ for the uncertainty Δr to be finite.

As such Δx or the uncertainty product becoming infinite does not violate the uncertainty relation (4), this however may not be desirable either. Notice that this ground state (10) is a much extended state that vanishes as $1/x$ asymptotically much slower as compared to the states in Fig. 1(a,b). Such a state, if useful in future, may be called a loosely bound and spatially extended state.

References

[1] Merzbacher E. 1970, *Quantum Mechanics* (John Wiley and Sons, Inc.: New York) 2^{nd}edition

[2] Griffith D. J. 2011 *Introduction to Quantum Mechanics* 2^{nd} edition (New-Delhi: Pearson)

[3] See p 84 in Ref. [2]

[4] Ahmed Z. and Yadav I. 2014 Position-momentum uncertainty products Eur. J. Phys. **35** 045015

[5] M. Ozawa 2003 Phys. Rev. A **67**, 042105; Jacqueline Erhart, Stephan Sponar, Georg Sulyok, Gerald Badurek, Masanao Ozawa & Yuji Hasegawa 2012 Nature Phys. **8** 185.

[6] Apostol T.M. 2002 *Calculus* (John Wiley and Sons, Inc.: New-York) 2^{nd}edition pp. 416-420

[7] See 2.5.4 on p 104 in Ref. [2]

[8] Abromowitz M., Stegun I.A 1970 *Handbook of Mathematical Functions* (Dover: New York) p. 255

[9] See pp 128-137 in Ref. [1]